Explicit solutions , conservation laws of the extended (2+1)-dimensional Jaulent-Miodek equation

Yu Li*, Xiqiang Liu and Xiangpeng Xin

School of Mathematical Sciences, Liaocheng University, Liaocheng 252059, People's Republic of China

arXiv:1512.09196v1 [nlin.SI] 31 Dec 2015

Abstract

By applying the direct symmetry method, the symmetry reductions and some new group invariant solutions were obtained, We have derived some exact solutions by using the relationship between the new solutions and the old ones, which include Weierstrass periodic solutions, elliptic periodic solutions, triangular function solutions and so on. Also, in order to reflect the characteristics and properties of this solutions, we give figures of some solutions. In addition, we give the conservation laws of the extended (2+1)-dimensional Jaulent-Miodek equation. At last, we draw conclusions and discuss it.

Keywords: Jaulent-Miodek equation, Symmetry, Exact solution.

1. Introduction

The nonlinear evolution equations(NLEEs) are encountered in a variety of scientific fields, such as chemistry, plasma physics, engineering , etc. In order to solve the problems of the nonlinear evolution equations, so many effective analytical and semianalytical approaches have been introduced, such as Lie point symmetry method[1–3], Exp-function method[4], tanh-function method[5, 6], Jacobi-elliptic method[7, 8], F-expansion method[9], Painleve analysis and the Hirota bilinear method[10, 11]. Especially, to find the Lie point symmetry and reduction of a given partial differential equation, the classical Lie group approach, the non-classical Lie group approach,the direct symmetry method have been suggested.

In this paper, we consider the extended (2+1)-dimensional Jaulent-Miodek equation. The Jaulent-Miodek equation refers to many branches of physics, such as condensed matter physics, fluid dynamics and optics. The relationship between Jaulent-Miodek equation and Euler-Darboux equation is found by Matsuno[12]. In recent years, many scientists have used many methods to obtain the exact solutions of Jaulent-Miodek equation, and have done a lot of work closely related to the spectral problem of Jaulent and Miodek[13, 14]. Fan[15]gives the Jaulent-Miodek equation with some traveling wave solutions, and Feng[16] gives the display solutions to the problem. For

* Corresponding author Email: liyu910118@sina.com

example, Exp-function method[4] is used to get the equation with periodic solution, tanh-coth method obtained the equation of soliton solutions, sign calculation equation is shaped as a cusp like singular soliton solutions, F -function equation of the traveling wave solutions are obtained[17–20].

The generaliged (2+1)-dimensional Jaulent-Miodek equation reads

$$u_{xt} - \alpha u_{xxxx} + \beta u_x^2 u_{xx} - u_{xx}u_y + \gamma u_x u_{xy} + \lambda u_{yy} + \delta u_{xx} = 0, \tag{1}$$

$\alpha, \beta, \gamma, \lambda$ and δ are real constants. The (2+1)-dimensional Jaulent-Miodek equation is generated by the (1+1)-dimensional Jaulent-Miodek hierarchy were developed[21]. When $\delta = 0$, a variety of solutions of the equation are obtained, please refer to the literature [22].

By applying the direct symmetry method, we get the symmetry reductions, group invariant solutions and some new exact solutions of Eq.(1), which include Weierstrass periodic solutions, Elliptic periodic solutions, Triangular function solutions and so on. This paper is organized in the following sections. In Sec.2. the Lie point symmetry groups and symmetry are obtain. And the brackets and the single parameter group are obtained, also the invariant solution of the group is obtained by the single parameter group. In Sec.3. we find the reduction and new solutions of the Eq.(1). In Sec.4. we give the conservation laws of Eq.(1). At last, we draw conclusions and discuss it.

2. Symmetry groups of the extended (2+1)-dimensional Jaulent-Miodek equation

By applying the classical Lie symmetry method we consider the one-parameter Lie group of infinitesimal transformations in (x, y, t, u) of Eq.(1) given by

$$\begin{aligned}
x^* &= x + \epsilon\xi(x, y, t, u) + \mathrm{O}\left(\epsilon^2\right), \\
y^* &= y + \epsilon\eta(x, y, t, u) + \mathrm{O}\left(\epsilon^2\right), \\
t^* &= t + \epsilon\tau(x, y, t, u) + \mathrm{O}\left(\epsilon^2\right), \\
u^* &= u + \epsilon\phi(x, y, t, u) + \mathrm{O}\left(\epsilon^2\right),
\end{aligned}$$

where ϵ is a small parameter. Thus, the vector field with the above group of transformations can be expressed as follows

$$V = \xi(x, y, t, u)\frac{\partial}{\partial x} + \eta(x, y, t, u)\frac{\partial}{\partial y} + \tau(x, y, t, u)\frac{\partial}{\partial t} + \phi(x, y, t, u)\frac{\partial}{\partial u}, \tag{2}$$

where the coefficient functions ξ, η, τ and ϕ are to be determined later.

If the vector field Eq.(1) generates a symmetry of Eq.(2) , then must satisfy Lie's symmetry condition

$$Pr^{(4)}V(\Delta)|_{\Delta=0} = 0, \tag{3}$$

where $\Delta = u_{xt} - \alpha u_{xxxx} + \beta u_x^2 u_{xx} - u_{xx}u_y + \gamma u_x u_{xy} + \lambda u_{yy} + \delta u_{xx}$. Applying the fourth prolongation $Pr^{(4)}V$ to Eq.(1), we find the following system of symmetry equations, and the invariant condition reads as

$$2\beta\phi^x u_x u_{xx}^2 + \gamma\phi^x u_{xy} + \phi^{xt} - \phi^{xx}u_y + \beta\phi^{xx}u_x^2 - \phi^y u_{xx} + \gamma\phi^{xy}u_x + \lambda\phi^{yy} + \delta\phi^{xx} - \alpha\phi^{xxxx} = 0, \tag{4}$$

where $\phi^x, \phi^{xx}, \phi^{xt}, \phi^{xy}, \phi^y, \phi^{yy}, \phi^{xxxx}$ are given explicitly in terms of ξ, η, τ, ϕ and the derivatives of ϕ. Furthermore, one can get

$$\begin{aligned}
\phi^x &= D_x(\phi - u_x\xi - u_t\tau - u_y\eta) + \xi u_{xx} + \eta u_{xy} + \tau u_{xt},\\
\phi^y &= D_y(\phi - u_x\xi - u_t\tau - u_y\eta) + \xi u_{xy} + \eta u_{yy} + \tau u_{yt},\\
\phi^{xx} &= D_{xx}(\phi - u_x\xi - u_t\tau - u_y\eta) + \xi u_{xxx} + \eta u_{xxy} + \tau u_{xxt},\\
\phi^{xt} &= D_{xt}(\phi - u_x\xi - u_t\tau - u_y\eta) + \xi u_{xxt} + \eta u_{xyt} + \tau u_{xtt},\\
\phi^{xy} &= D_{xy}(\phi - u_x\xi - u_t\tau - u_y\eta) + \xi u_{xxy} + \eta u_{xyy} + \tau u_{xyt},\\
\phi^{yy} &= D_{yy}(\phi - u_x\xi - u_t\tau - u_y\eta) + \xi u_{xyy} + \eta u_{yyy} + \tau u_{yyt},\\
\phi^{xxxx} &= D_{xxxx}(\phi - u_x\xi - u_t\tau - u_y\eta) + \xi u_{xxxxx} + \eta u_{xxxxy} + \tau u_{xxxxt},
\end{aligned} \tag{5}$$

where D_i denotes the total derivative operator and is defined as $D_i = \frac{\partial}{\partial x_i} + u_i\frac{\partial}{\partial u} + u_{ij}\frac{\partial}{\partial x_j} + \cdots$, and $(x_1, x_2, x_3) = (t, x, y)$.

By Lie symmetry analysis, setting the coefficients of the polynomial to zero, yields many differential equations about the function ξ, η, τ and ϕ, and solve the corresponding determined equations, one can get

$$\begin{aligned}
\xi &= \tfrac{c_1}{3}x - F_1(t) + \tfrac{2c_1\delta t}{3} + \tfrac{2\beta c_3 y}{-4\beta\lambda+\gamma^2-\gamma} + c_5,\\
\eta &= \tfrac{2c_1}{3}y + c_3 t + c_4,\\
\tau &= c_1 t + c_2,\\
\phi &= \tfrac{c_3(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} + F_1'(t)y + F_2(t),
\end{aligned} \tag{6}$$

where $c_i (i = 1, 2, 3, 4, 5)$ are arbitrary constants, $F_1(t)$ and $F_2(t)$ are arbitrary functions. Therefore, we obtain the following symmetries of Eq.(1),

$$\begin{aligned}
\sigma = &\left(\tfrac{c_1}{3}x - F_1(t) + \tfrac{2c_1\delta t}{3} + \tfrac{2\beta c_3 y}{-4\beta\lambda+\gamma^2-\gamma} + c_5\right)u_x + \left(\tfrac{2c_1}{3}y + c_3 t + c_4\right)u_y + (c_1 t + c_2)\,u_t\\
&- \left(\tfrac{c_3(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} + F_1'(t)\,y + F_2(t)\right).
\end{aligned} \tag{7}$$

From the above results, equivalent vector expression of the above symmetry is

$$\begin{aligned}
V = &\left(\tfrac{c_1}{3}x - F_1(t) + \tfrac{2c_1\delta t}{3} + \tfrac{2\beta c_3 y}{-4\beta\lambda+\gamma^2-\gamma} + c_5\right)\tfrac{\partial}{\partial x} + (c_1 t + c_2)\tfrac{\partial}{\partial t} + \left(\tfrac{2c_1}{3}y + c_3 t + c_4\right)\tfrac{\partial}{\partial y}\\
&+ \left(\tfrac{c_3(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} + F_1'(t)\,y + F_2(t)\right)\tfrac{\partial}{\partial u}.
\end{aligned} \tag{8}$$

Setting $F_1(t) = c_6 t + c_7$, $F_2(t) = c_8 t^2 + c_9 t + c_{10}$, where $c_i (i = 6, 7, 8, 9, 10)$ are arbitrary constants. From Eq.(8), the corresponding ten-dimensional Lie algebra = $\{V_1, V_2, V_3, V_4, V_5, V_6, V_7, V_8, V_9, V_{10}\}$ is reached, where

$$\begin{aligned}
&V_1 = (\tfrac{1}{3}x + \tfrac{2\delta t}{3})\tfrac{\partial}{\partial x} + \tfrac{2}{3}y\tfrac{\partial}{\partial y} + t\tfrac{\partial}{\partial t}, V_2 = \tfrac{\partial}{\partial t},\\
&V_3 = \tfrac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}\tfrac{\partial}{\partial x} + t\tfrac{\partial}{\partial y} + \tfrac{(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma}\tfrac{\partial}{\partial u},\\
&V_4 = \tfrac{\partial}{\partial y}, \quad V_5 = \tfrac{\partial}{\partial x}, \quad V_6 = -t\tfrac{\partial}{\partial x} + y\tfrac{\partial}{\partial u},\\
&V_7 = -\tfrac{\partial}{\partial x}, \quad V_8 = t^2\tfrac{\partial}{\partial u}, \quad V_9 = t\tfrac{\partial}{\partial u},\\
&V_{10} = \tfrac{\partial}{\partial u},
\end{aligned} \tag{9}$$

The one-parameter group generated by $v_i (i = 1, 2, \cdots, 10)$ is given as

$$\begin{aligned}
&g_1 : \left(t\delta e^{\varepsilon} + (-t\delta + x)\, e^{\frac{\varepsilon}{3}}, y e^{\frac{2}{3}\varepsilon}, t e^{\varepsilon}, u\right), g_2 : (x, y, t + \varepsilon, u), \\
&g_3 : \left(\int_0^{\varepsilon} \frac{2\beta(z_1 t + y)}{-4\beta\lambda + \gamma^2 - \gamma} dz_1 + x, y + t\varepsilon, t, \int_0^{\varepsilon} \frac{(\gamma - 1)\left(\int_0^{z_1} \frac{2\beta(z_1 t + y)}{-4\beta\lambda + \gamma^2 - \gamma} dz_1 + x\right)}{-4\beta\lambda + \gamma^2 - \gamma} dz_1 + u \right), \\
&g_4 : (x, y + \varepsilon, t, u), g_5 : (\varepsilon + x, y, t, u), g_6 : (x - t\varepsilon, y, t, y\varepsilon + u), \\
&g_7 : (-\varepsilon + x, y, t, u), g_8 : \left(x, y, t, t^2\varepsilon + u\right), g_9 : (x, y, t, t\varepsilon + u), \\
&g_{10} : (x, y, t, \varepsilon + u).
\end{aligned} \tag{10}$$

From the above groups $g_i (i = 1, 2, \cdots, 10)$, one can get the following solutions of Eq.(1)

$$\begin{aligned}
&u_1 = f\left(-\frac{\delta t e^{\varepsilon} - \delta t e^{\frac{\varepsilon}{3}} - x e^{\varepsilon}}{e^{\frac{\varepsilon}{3}} e^{\varepsilon}}, y e^{-\frac{2}{3}\varepsilon}, t e^{-\varepsilon}\right), u_2 = f(x, y, t - \varepsilon), \\
&u_3 = f\left(x - \int_0^{\varepsilon} \frac{2\beta(z_1 t - t\varepsilon + y)}{-4\beta\lambda + \gamma^2 - \gamma} dz_1, y - t\varepsilon, t\right) + \int_0^{\varepsilon} \frac{(\gamma - 1)\left(\int_0^{z_1} \frac{2\beta(z_1 t + y)}{-4\beta\lambda + \gamma^2 - \gamma} dz_1 + x\right)}{-4\beta\lambda + \gamma^2 - \gamma} dz_1, \\
&u_4 = f(x, y - \varepsilon, t), u_5 = f(x - \varepsilon, y, t), u_6 = f(x + t\varepsilon, y, t) + y\varepsilon, \\
&u_7 = f(x + \varepsilon, y, t), u_8 = f(x, y, t) + t^2\varepsilon, u_9 = f(x, y, t) + t\varepsilon, \\
&u_{10} = f(x, y, t) + \varepsilon,
\end{aligned} \tag{11}$$

where ϵ is a parameter , f is an arbitrary known solution of Eq.(1).

3. Similarity reductions and new exact solutions of the extended (2+1)-dimensional Jaulent-Miodek equation

To search for more similarity reductions of (1), one should consider Lie point symmetries of (1).In this section, we will discuss how to construct similarity reductions and solutions of Eq.(1). The characteristic equations have been written from Eq.(7) as follows

$$\frac{dx}{\frac{c_1}{3}x - F_1(t) + \frac{2c_1\delta t}{3} + \frac{2\beta c_3 y}{-4\beta\lambda + \gamma^2 - \gamma} + c_5} = \frac{dy}{\frac{2c_1}{3}y + c_3 t + c_4} = \frac{dt}{c_1 t + c_2} = \frac{du}{\frac{c_3(\gamma - 1)x}{-4\beta\lambda + \gamma^2 - \gamma} + F_1'(t)\, y + F_2(t)}. \tag{12}$$

Here we discuss the following Cases.

Case 1 Letting $c_2 = c_4 = 1, c_1 = c_3 = c_5 = 0, F_1(t) = 0, F_2(t) \neq 0$, solving the corresponding characteristic equations, we have the following invariants ξ, η and the function u

$$\xi = x, \eta = t - y, \qquad u = \int F_2(t)\, dt + f(\xi, \eta), \tag{13}$$

Substituting Eq.(13) into Eq.(1) , we have

$$f_{\xi\eta} - \alpha f_{\xi\xi\xi\xi} + \beta f_{\xi}^2 f_{\xi\xi} + f_{\xi\xi} f_{\eta} - \gamma f_{\xi} f_{\xi\eta} + \lambda f_{\eta\eta} + \delta f_{\xi\xi} = 0, \tag{14}$$

In order to get the exact solution of Eq.(14), we look for the Lie point transformation group of Eq.(14), and the

corresponding vector field of Eq.(14) as follows

$$\widetilde{V_1} = \left(\frac{a}{2}\xi + \frac{a\eta\beta}{4\beta\lambda - \gamma^2 + \gamma} + d\right)\frac{\partial}{\partial\xi} + (a\eta + b)\frac{\partial}{\partial\eta} + \left(-\frac{a(\gamma-1)\xi}{8\beta\lambda - 2\gamma^2 + 2\gamma} - \frac{a\left(-4\beta\delta\lambda + \delta\gamma^2 + \beta - \delta\gamma\right)\eta}{-4\beta\lambda + \gamma^2 - \gamma} + c\right)\frac{\partial}{\partial f}, \quad (15)$$

where a, b, c and d are arbitrary constants.

Furthermore, we can get the characteristic equation of Eq.(14),

$$\frac{d\xi}{\frac{a}{2}\xi + \frac{a\eta\beta}{4\beta\lambda-\gamma^2+\gamma} + d} = \frac{d\eta}{a\eta + b} = \frac{df}{-\frac{a(\gamma-1)\xi}{8\beta\lambda-2\gamma^2+2\gamma} - \frac{a\left(-4\beta\delta\lambda+\delta\gamma^2+\beta-\delta\gamma\right)\eta}{-4\beta\lambda+\gamma^2-\gamma} + c}, \quad (16)$$

where a, b, c and d are arbitrary constants.

Case 1.1 $a = 0, c = 0, b \neq 0, d \neq 0$

In this case, we can get the following solution of Eq.(14)

$$\theta = b\xi - d\eta, \qquad f = g(\theta), \quad (17)$$

and $g(\theta)$ need satisfy

$$\left(\lambda d^2 + \delta b^2 - bd\right)g'' - \alpha b^4 g^{(4)} + \beta b^4 g'^2 g'' + \left(\gamma b^2 d - b^2 d\right)g'g'' = 0, \quad (18)$$

Integral to Eq.(18) , yields

$$\left(\lambda d^2 + \delta b^2 - bd\right)g' - \alpha b^4 g''' + \frac{1}{3}\beta b^4 g'^3 + \frac{1}{2}\left(\gamma b^2 d - b^2 d\right)g'^2 + C_1 = 0, \quad (19)$$

where D is an arbitrary constant.

Letting $k_1 = \lambda d^2 + \delta b^2 - bd, k_2 = \frac{1}{3}\beta b^4, k_3 = \frac{1}{2}\left(\gamma b^2 d - b^2 d\right), k_4 = \alpha b^4$, Eq.(19) can be reads as

$$k_1 g' + k_2 g'^3 + k_3 g'^2 - k_4 g''' + C_1 = 0, \quad (20)$$

Setting $g'(\xi) = p$, Eq.(20) can be further simplified to

$$k_1 p + k_2 p^3 + k_3 p^2 - k_4 p'' + C_1 = 0, \quad (21)$$

In order to get the solutions of Eq.(21) , We adopt two methods. The generalized tanh function expansion method[24] to be adopted in first one . There are explicit solutions to the following form

$$p = \sum_{i=0}^{n} q_i Y(\theta)^i + \sum_{i=1}^{n} q_{-i} Y(\theta)^{-i}, \quad (22)$$

where $\theta = b\xi - d\eta$, and n, q_i, q_{-i} are undetermined constants, also Y is the solution of the *Riccati* equation as

follows

$$Y'(\theta) = A + BY(\theta) + CY^2(\theta), \tag{23}$$

where A, B and C are constants.

Balancing the highest order derivative term and nonlinear term in Eq.(21) , it follows that $n = 1$. Therefore the explicit solution of the equation can be expressed as

$$p = q_0 + q_1 Y(\theta) + \frac{q_{-1}}{Y(\theta)}, \tag{24}$$

By substituting Eq.(23) and Eq.(24) into Eq.(21) , we can obtain the following solutions

$$\begin{aligned} p_1 &= R + \sqrt{\tfrac{2k_4}{k_2}} C [\sec(\theta) - \tan(\theta)] + \sqrt{\tfrac{2k_4}{k_2}} A [\sec(\theta) - \tan(\theta)]^{-1}, \\ p_2 &= R + \sqrt{\tfrac{2k_4}{k_2}} C \tfrac{\tan\theta}{1 \pm \sec\theta} + \sqrt{\tfrac{2k_4}{k_2}} A \left(\tfrac{\tan\theta}{1 \pm \sec\theta}\right)^{-1}, \\ p_3 &= R + \sqrt{\tfrac{2k_4}{k_2}} C \tfrac{\exp(B\theta) - A}{B} + \sqrt{\tfrac{2k_4}{k_2}} A \left[\tfrac{\exp(B\theta) - A}{B}\right]^{-1}, \end{aligned} \tag{25}$$

where $\theta = b\xi - d\eta$ and $R = \frac{-k_3 + \sqrt{k_3^2 - 12k_2k_4AC - 3k_1k_2 + 3k_2k_4B^2}}{3k_2}$.

The solutions of the original equation are obtained, respectively,

$$\begin{aligned} u_1 &= \int F_2(t)\,dt + R\theta + \sqrt{\tfrac{2k_4}{k_2}} C [\ln(\sec(\theta) + \tan(\theta)) + \ln(\cos(\theta))] - \sqrt{\tfrac{2k_4}{k_2}} A \ln(\sin(\theta) - 1) + C_2, \\ u_2 &= \int F_2(t)\,dt + R\theta - \sqrt{\tfrac{2k_4}{k_2}} C [\ln(\sec(\theta) \pm 1) - \ln(\sec(\theta))] + \sqrt{\tfrac{2k_4}{k_2}} A [\ln(\sin(\theta)) \pm \ln(\csc(\theta) - \cot(\theta))] + C_3, \\ u_3 &= \int F_2(t)\,dt + R\theta - \sqrt{\tfrac{2k_4}{k_2}} C \tfrac{e^{(B\theta)}}{B} - A\theta - \sqrt{\tfrac{2k_4}{k_2}} A \left[\tfrac{\ln(-e^{(B\theta)} + A)}{A} - \tfrac{\ln(e^{(B\theta)})}{A}\right] + C_4, \end{aligned} \tag{26}$$

where C_1, C_2, C_3 are integral constants. When $\delta = 0$, we get the Painleve II solutions, but we get the other new exact solutions from this article.

The second, by applying Jacobi-elliptic function expansion method, we look for the solitary wave solutions of Eq.(21). Suppose the solutions has the following form

$$G = p(\theta) = \sum_{i=0}^{n} a_i sn^i(\theta, m) + \sum_{i=0}^{n} b_i sn^{-i}(\theta, m). \tag{27}$$

Balancing the highest order derivative term and nonlinear term in Eq.(21), it follows that $n = 1$. Therefore, the explicit solution of Eq.(27) can be expressed as

$$p(\theta) = a_0 + a_1 sn(\theta, m) + b_1 sn^{-1}(\theta, m), \tag{28}$$

where a_0, a_1, b_1 are undetermined constants. In the same way, we can obtain the following solutions

$$\begin{aligned} G_1 &= p_4 = \frac{-k_3 + \sqrt{k_3^2 - 18k_2k_4m - 3k_1k_2 - 3k_2k_4 - 3k_2k_4m^2}}{3k_2} - \sqrt{\tfrac{2k_4}{k_2}} \left[sn(\theta, m) - sn^{-1}(\theta, m)\right], \\ G_2 &= p_5 = \frac{-k_3 + \sqrt{k_3^2 - 3k_2k_4 - 3k_1k_2 - 3k_2k_4m^2}}{3k_2} - \sqrt{\tfrac{2k_4}{k_2}} m sn(\theta, m), \end{aligned} \tag{29}$$

where $0 < m < 1$, m is modulus of elliptic functions.

So we can get the other set of solutions for Eq.(1)

$$\begin{aligned} u_4 &= \int F_2(t)\,dt + \frac{-k_3+\sqrt{k_3^2-18k_2k_4m-3k_1k_2-3k_2k_4-3k_2k_4m^2}}{3k_2}\theta - \sqrt{\frac{2k_4}{k_2}}\left[\frac{1}{m}\ln\left(dn(\theta,m)-mcn(\theta,m)\right)-\ln\left(\frac{sn(\theta,m)}{cn(\theta,m)+dn(\theta,m)}\right)\right], \\ u_5 &= \int F_2(t)\,dt + \frac{-k_3+\sqrt{k_3^2-3k_2k_4-3k_1k_2-3k_2k_4m^2}}{3k_2}\theta - \sqrt{\frac{2k_4}{k_2}}\ln\left(dn(\theta,m)-mcn(\theta,m)\right). \end{aligned} \tag{30}$$

Case 1.2 $a=0, d\neq 0, b\neq 0, c\neq 0$

In this case, we can get the following solution of Eq.(14) ,

$$\omega = \frac{c\eta}{b}, \theta = \xi, \qquad f = \omega + h(\theta), \tag{31}$$

Take Eq.(31) into Eq.(14), we can get

$$-\alpha h^{(4)} + \beta h'^2 h'' + \left(\frac{c}{b}+\delta\right)h'' = 0, \tag{32}$$

Setting $h'(\xi) = W$, and Eq.(32) can be written as

$$-\alpha W'' + \frac{\beta}{3}W^3 + \left(\frac{c}{b}+\delta\right)W + C_5 = 0, \tag{33}$$

Letting $A = \frac{\beta}{3\alpha}, B = \frac{c}{b\alpha}+\frac{\delta}{\alpha}, r = \frac{C_5}{\alpha}$, then Eq.(33) can be written as

$$W'' = AW^3 + BW + R, \tag{34}$$

Taking different values of A and B , and Eq.(34) also have the corresponding different solutions,

$$\begin{aligned} &(1)\,If\,A<0, B>0, R=0, V_1 = \sqrt{\frac{-B}{A}}\,\mathrm{sec}\,h\left(\sqrt{B}\theta\right), \\ &(2)\,If\,A>0, B<0, R=0, V_2 = \sqrt{\frac{-B}{A}}\,\mathrm{sec}\,h\left(\sqrt{-B}\theta\right), \\ &(3)\,If\,A>0, B>0, R=\frac{B^2}{4A}, V_3 = \sqrt{\frac{B}{2A}}\,\mathrm{sec}\,h\left(\sqrt{B/2}\theta\right), \\ &(4)\,If\,A=\frac{4}{m}, B=-6m-m^2-1, R=2m^3+m^4+m^2, V_4 = \frac{mdn(\theta,m)cn(\theta,m)}{msn^2(\theta,m)-1}, \\ &(5)\,If\,A=-4m_1, B=-6m_1-m^2+2, R=2-2m_1-m^2, V_5 = -\frac{m^2sn(\theta,m)cn(\theta,m)}{m_1+dn^2(\theta,m)}, \\ &(6)\,If\,A=m^2\left(-1+m^2\right), B=2m^2-1, R=1, V_6 = sd(\theta) = \frac{sn(\theta,m)}{dn(\theta,m)}, \\ &(7)\,If\,A=1-m^2, B=2m^2-1, R=-m^2, V_7 = nc(\theta,m) = \frac{1}{cn(\theta,m)}, \\ &(8)\,If\,A=1, B=2-m^2, R=1, V_8 = \frac{sn(\theta,m)dn(\theta,m)}{cn(\theta,m)}, \\ &(9)\,If\,A=\frac{1}{4}, B=\frac{m^2-2}{2}, R=\frac{m^2}{4}, V_9 = ns(\theta,m)+ds(\theta,m), \\ &(10)\,If\,A=1, B=2+m^2, R=-2m^2+m^4+1, V_{10} = \frac{dn(\theta,m)cn(\theta,m)}{sn(\theta,m)}, \end{aligned} \tag{35}$$

where m is modulus of elliptic functions.

Applying the above expressions of explicit solutions $W_i\,(i=1,\ldots 10)$, one can obtain h and $f=\omega+h(\theta)$, and

we can derive the corresponding solutions of Eq.(1). For example

$$\begin{aligned} u_6 &= \tfrac{c\eta}{b} + \sqrt{\tfrac{-1}{A}} \arctan\left(\sinh\left(\sqrt{B}\theta\right)\right), \\ u_7 &= \tfrac{c\eta}{b} - \ln\left(sn\left(\theta, m\right)\right), \\ u_8 &= \tfrac{c\eta}{b} + \ln\left(ds\left(\theta, m\right) - cs\left(\theta, m\right)\right) + \ln\left(ns\left(\theta, m\right) - cs\left(\theta, m\right)\right). \end{aligned} \tag{36}$$

Case 2 Letting $c_2 = 1, c_1 = c_3 = c_4 = c_5 = 0, F_1(t) = 0, F_2(t) \neq 0$, in the same way, we can get

$$\xi = x, \eta = y, \qquad u = \int F_2(t)\, dt + f(\xi, \eta), \tag{37}$$

Substituting Eq.(37) into Eq.(1) , we have

$$-\alpha f_{\xi\xi\xi\xi} + \beta f_{\xi}^2 f_{\xi\xi} - f_{\xi\xi} f_{\eta} + \gamma f_{\xi} f_{\xi\eta} + \lambda f_{\eta\eta} + \delta f_{\xi\xi} = 0. \tag{38}$$

Making transformation with $f(\xi, \eta) = f(w), w = l\xi + k\eta$, we can get

$$-\alpha l^4 f^{(4)} + \beta l^4 f'^2 f'' - l^2 k f' f'' + \gamma l^2 k f' f'' + \lambda k^2 f'' + \delta l^2 f'' = 0. \tag{39}$$

Integrating twice with respect to f , yields

$$f''^2 = \frac{\beta}{6\alpha} f'^4 + \frac{(\gamma - 1)k}{\alpha l^2} f'^3 + \frac{\lambda k^2 + \delta l^2}{\alpha l^4} f'^2 + \frac{2C_6}{\alpha l^4} f' + \frac{2C_7}{\alpha l^4} = 0, \tag{40}$$

where d_0, d_1 are arbitrary constants. Letting $f' = H$, and Eq.(40) can be written as

$$H' = \mu \sqrt{h_0 + h_1 H + h_2 H^2 + h_3 H^3 + h_4 H^4}, \tag{41}$$

where $f = f(w), \mu = \pm 1$, and

$$h_0 = \frac{2C_7}{\alpha l^4}, h_1 = \frac{2C_6}{\alpha l^4}, h_2 = \frac{\lambda k^2 + \delta l^2}{\alpha l^4}, h_3 = \frac{(\gamma - 1)k}{\alpha l^2}, h_4 = \frac{\beta}{6\alpha}. \tag{42}$$

We can get the following solutions through the following discussion. The explicit solution of Eq.(41) have been given in [22, 23].

Case 2.1 Polynomial solutions

If $h_0 > 0, h_1 = h_2 = h_3 = h_4 = 0$, then $H = \mu \sqrt{h_0} w + C_8$, and we obtain

$$u_9 = \int F_2(t)\, dt + \frac{1}{2}\mu \sqrt{h_0} w^2 + C_8 w. \tag{43}$$

If $h_1 \neq 0, h_2 = h_3 = h_4 = 0$, then $H = -\frac{h_0}{h_1} + \mu^2 \left(\frac{h_1}{4}\right) w^2 + C_9$, then the solution of Eq.(1) is expressed by

$$u_{10} = \int F_2(t)\, dt + \frac{1}{3}\mu^2 \frac{h_1}{4} w^3 + \left(C_9 - \frac{h_0}{h_1}\right) w. \tag{44}$$

where $w = l\xi + k\eta$, and ξ, η are arbitrary constants.

Case 2.2 Rational solutions

If $h_4 > 0, h_0 = h_1 = h_2 = h_3 = 0$, then $H = -\frac{1}{\mu\sqrt{h_4}w} + C_{10}$, then we obtain

$$u_{11} = \int F_2(t)\,dt + \frac{\ln w}{\mu\sqrt{h_4}} + C_{10}w. \tag{45}$$

If $h_2 > 0, h_0 = h_1 = h_3 = h_4 = 0$, then $H = e^{\mu\sqrt{h_2}w} + C_{11}$, and one can derive

$$u_{12} = \int F_2(t)\,dt + \frac{2\sqrt{w}e^{\mu\sqrt{h_2}w}}{\mu\sqrt{h_2}} + C_{11}w. \tag{46}$$

where $w = l\xi + k\eta$, and ξ, η are arbitrary constants.

Case 2.3 Triangular function solutions

(1) If $h_0 = h_1 = 0$, then we can get

$$\begin{aligned} H_1(w) &= \frac{4h_2C_0\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}{2\mu(1-\Delta_1)+2\mu(1+\Delta_1)\tanh\left(\frac{\sqrt{h_2}}{2}w\right)-(2h_3C_0+1-\Delta_1)\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}, \\ H_2(w) &= \frac{4h_2C_0\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}{2\mu\Theta-2\mu\left(\Delta_2+C_0^2\right)\tanh\left(\frac{\sqrt{h_2}}{2}w\right)-(2h_3C_0+\Theta)\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}, \end{aligned} \tag{47}$$

where the constants $h_2 > 0$, $C_0 = \exp\left(\sqrt{h_2}C_{12}\right), \Delta_1 = C_0^2\left(4h_2h_4 - h_3^2\right), \Delta_2 = 4h_2h_4 - h_3^2$ and $\Theta = C_0^2 - \Delta_2$.

Then the solution of Eq.(1) is expressed by

$$\begin{aligned} u_{13} &= \int F_2(t)\,dt + \int \frac{4h_2C_0\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}{2\mu(1-\Delta_1)+2\mu(1+\Delta_1)\tanh\left(\frac{\sqrt{h_2}}{2}w\right)-(2h_3C_0+1-\Delta_1)\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}dw + C_{13}, \\ u_{14} &= \int F_2(t)\,dt + \int \frac{4h_2C_0\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}{2\mu\Theta-2\mu\left(\Delta_2+C_0^2\right)\tanh\left(\frac{\sqrt{h_2}}{2}w\right)-(2h_3C_0+\Theta)\sec h^2\left(\frac{\sqrt{h_2}}{2}w\right)}dw + C_{14}. \end{aligned} \tag{48}$$

(2) If $h_1 = h_3 = 0, h_0 = \frac{h_2^2}{4h_4}, h_2 > 0, h_4 > 0$, then $H = \mu\sqrt{\frac{h_2}{h_4}}\tan\left(\sqrt{\frac{h_2}{2}}w\right)$, we can get

$$u_{15} = \int F_2(t)\,dt + \mu\sqrt{\frac{1}{2h_4}}\ln\left(1 + \tan^2\left(\sqrt{\frac{h_2}{2}}w\right)\right) + C_{15}. \tag{49}$$

where $w = l\xi + k\eta$, and ξ, η are arbitrary constants.

In order to reflect the characteristics and properties of the solution u_{15}, we give the image of the solution u_{15}, as follows

Case 2.4 Elliptic periodic solutions

If $h_1 = h_3 = 0$, then we can obtain some new Jacobi-elliptic function solutions

(1)

$$H_3 = \mu\sqrt{\frac{-h_2m^2}{h_4\left(2m^2-1\right)}}cn\left(\sqrt{\frac{h_2}{2m^2-1}}w, m\right), h_4 < 0, h_2 > 0, h_0 = \frac{h_2^2m^2\left(1-m^2\right)}{h_4\left(2m^2-1\right)^2}. \tag{50}$$

So Eq.(1) has the following solution

$$u_{16} = \int F_2(t)\,dt + \mu\sqrt{\frac{-h_2m^2}{h_4(2m^2-1)}}\int cn\left(\sqrt{\frac{h_2}{2m^2-1}}w, m\right)dw + C_{16}. \tag{51}$$

(2)

$$H_4(w) = \mu\sqrt{\frac{-h_2}{h_4(2-m^2)}}dn\left(\sqrt{\frac{h_2}{2-m^2}}w, m\right), h_4 < 0, h_2 > 0, h_0 = \frac{h_2^2\left(1-m^2\right)}{h_4(2-m^2)^2}, \tag{52}$$

And one can derive

$$u_{17} = \int F_2(t)\,dt + \mu\sqrt{\frac{-h_2}{h_4(2-m^2)}}\int dn\left(\sqrt{\frac{h_2}{2-m^2}}w, m\right)dw + C_{17}. \tag{53}$$

(3)

$$H_5(w) = \mu\sqrt{\frac{-h_2m^2}{h_4(m^2+1)}}sn\left(\sqrt{-\frac{h_2}{m^2+1}}w, m\right), h_4 > 0, h_2 < 0, h_0 = \frac{h_2^2m^2}{h_4(m^2+1)^2}, \tag{54}$$

And we can obtain

$$u_{18} = \int F_2(t)\,dt + \mu\sqrt{\frac{-h_2m^2}{h_4(m^2+1)}}\int sn\left(\sqrt{-\frac{h_2}{m^2+1}}w, m\right)dw + C_{18}. \tag{55}$$

where $0 < m < 1$, m is modulus of elliptic functions, $w = l\xi + k\eta$, and ξ, η are arbitrary constants.

When $m \to 1$ and $m \to 0$The Jacobi elliptic function is degenerated into trigonometric or hyperbolic functionas follows

$m \to 1, sn(w,m) \to \tanh(w), cn(w,m) \to \sec h(w), dn(w,m) \to \sec h(w),$

$m \to 0, sn(w,m) \to \sin(w), cn(w,m) \to \cos(w), dn(w,m) \to 1.$

Case 2.5 Weierstrass periodic solutions

If $h_2 = h_4 = 0$, then Eq.(41) can get some Weierstrass periodic solutions

$H_6(w) = \mu\wp\left(\frac{\sqrt{h_3}}{2}w, g_2, g_3\right)$, where $h_3 > 0, g_2 = -4\frac{h_1}{h_3}, g_3 = -4\frac{h_0}{h_3}$.

So we can derive solution of Eq.(1)

$$u_{19} = \int F_2(t)\,dt + \mu\int\wp\left(\frac{\sqrt{h_3}}{2}w, g_2, g_3\right)dw + C_{19}. \tag{56}$$

Case 3 Letting $c_2 = c_4 = 1, c_1 = c_3 = c_5 = 0, F_1(t) = a, F_2(t) = b$, solving the corresponding characteristic equations, we have the following invariants ξ, η and the function u .

$$\xi = y + \tfrac{x}{a}, \eta = t + \tfrac{x}{a}, \qquad u = -\tfrac{b}{a}x + f(\xi, \eta), \tag{57}$$

Substituting Eq.(57) into Eq.(1) , we have

$$\begin{aligned}&\left(\tfrac{\lambda a^2+\delta-b\gamma}{a^2}+\tfrac{b^2\beta}{a^4}\right)f_{\xi\xi}+\left(\tfrac{2a-b\gamma+2\delta}{a^2}+\tfrac{2b^2\beta}{a^4}\right)f_{\xi\eta}+\left(\tfrac{a+\delta}{a^2}+\tfrac{b^2\beta}{a^4}\right)f_{\eta\eta}+\left(\tfrac{\gamma-1}{a^2}-\tfrac{2b\beta}{a^4}\right)f_\xi f_{\xi\xi}\\&+\left(\tfrac{\gamma-2}{a^2}-\tfrac{4b\beta}{a^4}\right)f_\xi f_{\xi\eta}-\left(\tfrac{1}{a^2}+\tfrac{2b\beta}{a^4}\right)f_\xi f_{\eta\eta}+\left(\tfrac{\gamma}{a^2}-\tfrac{4b\beta}{a^4}\right)f_\eta f_{\xi\eta}+\left(\tfrac{\gamma}{a^2}-\tfrac{2b\beta}{a^4}\right)f_\eta f_{\xi\xi}-\tfrac{4b\beta}{a^4}f_\eta f_{\eta\eta}\\&-\tfrac{\alpha}{a^4}\left(f_{\xi\xi\xi\xi}+4f_{\xi\xi\xi\eta}+6f_{\xi\xi\eta\eta}+4f_{\xi\eta\eta\eta}+f_{\eta\eta\eta\eta}\right)+\tfrac{\beta}{a^4}\left(f_\xi+f_\eta\right)^2\left(f_{\xi\xi}+f_{\eta\eta}+2f_{\xi\eta}\right)=0.\end{aligned}\tag{58}$$

Making transformation with $f(\xi,\eta)=f(\Omega),\Omega=k\xi+l\eta$,we can get

$$\begin{aligned}&\left(\tfrac{kl(2a-b\gamma+2\delta)+(a+\delta)l^2+\left(\lambda a^2+\delta-b\gamma\right)k^2}{a^2}+\tfrac{b^2\beta(l+k)^2}{a^4}\right)f''+\left(\tfrac{(\gamma-1)\left(k^3+2lk^2+l^2k\right)}{a^2}-\tfrac{2b\beta(k+l)^3}{a^4}\right)f'f''\\&+\tfrac{\beta(l+k)^4}{a^4}f'^2f''-\tfrac{\alpha(l+k)^4}{a^4}f^{(4)}=0.\end{aligned}\tag{59}$$

Integrating twice with respect to f ,yields

$$\begin{aligned}f''^2=&\tfrac{\beta}{6\alpha}f'^4+\left(\tfrac{a^2(\gamma-1)\left(k^3+2lk^2+l^2k\right)}{3\alpha(l+k)^4}-\tfrac{2b\beta}{3\alpha(l+k)}\right)f'^3\\&+\left(\tfrac{a^2\left[kl(2a-b\gamma+2\delta)+(a+\delta)l^2+\left(\lambda a^2+\delta-b\gamma\right)k^2\right]}{\alpha(l+k)^4}+\tfrac{2b^2\beta}{\alpha(l+k)^2}\right)f'^2\\&+\tfrac{2a^4e_0}{\alpha(l+k)^4}f'+\tfrac{2a^4e_1}{\alpha(l+k)^4}.\end{aligned}\tag{60}$$

Eq.(40) and Eq.(60) are all elliptic functions, so the solutions of case 3 is reference to case 2 .

Case 4 Letting $c_4=1,c_1=c_2=c_3=c_5=0,F_1(t)=0,F_2(t)=t^2$, in the same way, one can derive

$$\xi=x,\eta=t,\qquad u=t^2y+f(\xi,\eta),\tag{61}$$

Substituting Eq.(61) into Eq.(1), we get

$$f_{\xi\eta}-\alpha f_{\xi\xi\xi\xi}+\beta f_\xi^2f_{\xi\xi}-\eta^2f_{\xi\xi}+\delta f_{\xi\xi}=0.\tag{62}$$

Assuming Eq.(62) has the following solution

$$f=s(\eta)\xi+z(\eta),\tag{63}$$

where $s(\eta)$ and $z(\eta)$ are the functions to be determined. Substituting Eq.(63) into Eq.(62) yields

$$f=Z_1\xi+z(t).\tag{64}$$

Combining Eq.(61) and Eq.(64) , then the new exact solution of the Eq.(1)

$$u_{20}=t^2y+Z_1x+z(t).\tag{65}$$

where Z_1 is integral constant.

4. Conservation laws

In this section, we will study the conservation laws by using the adjoint equation and symmetries of Eq.(1). For Eq.(1), the adjoint equation has the form

$$v_{xt} - \alpha v_{xxxx} + \beta u_x^2 v_{xx} - 2\beta u_{xx} u_x v_x - v_{xx} u_y + u_{xx} v_y - \gamma v_x u_{xy} + \gamma u_x v_{xy} + \lambda v_{yy} + \delta v_{xx} = 0. \tag{66}$$

and the Lagrangian in the symmetrized form

$$L = v\left(u_{xt} - \alpha u_{xxxx} + \beta u_x^2 u_{xx} - u_{xx} u_y + \gamma u_x u_{xy} + \lambda u_{yy} + \delta u_{xx}\right). \tag{67}$$

Theorem 1. Every Lie point, Lie-Bäclund and non-local symmetry of Eq.(1) provides a conservation law for Eq.(1) and the adjoint equation. Then the elements of conservation vector (C^1, C^2, C^3) are defined by the following expression:

$$\begin{aligned} C^i = {} & \xi^i L + W^\alpha \left[\frac{\partial L}{\partial u^\alpha_{ij}} - D_j\left(\frac{\partial L}{\partial u^\alpha_{ij}}\right) + D_j D_k\left(\frac{\partial L}{\partial u^\alpha_{ijk}}\right) - D_j D_k D_r\left(\frac{\partial L}{\partial u^\alpha_{ijkr}}\right) + \cdots\right] \\ & + D_j(W)\left[\frac{\partial L}{\partial u^\alpha_{ij}} - D_k\left(\frac{\partial L}{\partial u^\alpha_{ijk}}\right) + \cdots\right] \\ & + D_j D_k(W)\left[\frac{\partial L}{\partial u^\alpha_{ijk}} - D_r\left(\frac{\partial L}{\partial u^\alpha_{ijkr}}\right) + \cdots\right], \end{aligned} \tag{68}$$

where W is the Lie characteristic function and

$$W^\alpha = \eta^\alpha_* - \xi^j u^\alpha_j. \tag{69}$$

The conserved vector corresponding to an operator is

$$v = \xi^1 \frac{\partial}{\partial t} + \xi^2 \frac{\partial}{\partial x} + \xi^3 \frac{\partial}{\partial y} + \xi^4 \frac{\partial}{\partial u}. \tag{70}$$

The operator v yields the conservation laws $D_t\left(C^1\right) + D_x\left(C^2\right) + D_y\left(C^3\right) = 0$, where the conserved vector $C = \left(C^1, C^2, C^3\right)$ is given by Eq.(68) and has the components

$$\begin{aligned} C^1 = {} & \xi^1 L + W\left(\frac{\partial L}{\partial u_t} - D_x\left(\frac{\partial L}{\partial u_{xt}}\right)\right) + W_x\left(\frac{\partial L}{\partial u_{xt}}\right), \\ C^2 = {} & \xi^2 L + W\left(\frac{\partial L}{\partial u_x} - D_x\left(\frac{\partial L}{\partial u_{xx}}\right) - D_y\left(\frac{\partial L}{\partial u_{xy}}\right) - D_t\left(\frac{\partial L}{\partial u_{xt}}\right) - D_{xxx}\left(\frac{\partial L}{\partial u_{xxxx}}\right)\right) \\ & + W_x\left(\frac{\partial L}{\partial u_{xx}} + D_{xx}\left(\frac{\partial L}{\partial u_{xxxx}}\right)\right) + W_{xx}\left(-D_x\left(\frac{\partial L}{\partial u_{xxxx}}\right)\right) + W_{xxx}\left(\frac{\partial L}{\partial u_{xxxx}}\right) \\ C^3 = {} & \xi^3 L + W\left(\frac{\partial L}{\partial u_y} - D_x\left(\frac{\partial L}{\partial u_{xy}}\right) - D_y\left(\frac{\partial L}{\partial u_{yy}}\right)\right) + W_x\left(\frac{\partial L}{\partial u_{xy}}\right) + W_y\left(\frac{\partial L}{\partial u_{yy}}\right). \end{aligned} \tag{71}$$

So Eq.(71) defined the corresponding components of non-local conservation law for the system of Eq.(1) and Eq.(66) corresponding to any operator admitted by Eq.(1).

Let us make conclusions for the operator

$$\begin{aligned} V = {} & \left(\tfrac{c_1}{3}x - F_1(t) + \tfrac{2c_1\delta t}{3} + \tfrac{2\beta c_3 y}{-4\beta\lambda+\gamma^2-\gamma} + c_5\right)\tfrac{\partial}{\partial x} + (c_1 t + c_2)\tfrac{\partial}{\partial t} + \left(\tfrac{2c_1}{3}y + c_3 t + c_4\right)\tfrac{\partial}{\partial y} \\ & + \left(\tfrac{c_3(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} + F_1'(t)y + F_2(t)\right)\tfrac{\partial}{\partial u}. \end{aligned} \tag{72}$$

For this operator, one can get

$$W = \frac{(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} - \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_x - tu_y. \tag{73}$$

We can get the conservation vector of Eq.(1)

$$\begin{aligned}
C^1 &= -\left(\frac{(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} - \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_x - tu_y\right)v_x + \left(\frac{(\gamma-1)}{-4\beta\lambda+\gamma^2-\gamma} - \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_{xx} - tu_{xy}\right)v,\\
C^2 &= -\frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}v\left(u_{xt} - \alpha u_{xxxx} + \beta u_x^2 u_{xx} - u_{xx}u_y + \gamma u_x u_{xy} + \lambda u_{yy} + \delta u_{xx}\right)\\
&\quad + \left(\frac{(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} - \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_x - tu_y\right)\left(u_{xy}v + \beta v_x u_x^2 - v_x u_y + \delta v_x - \gamma v_y u_x - v_t + \alpha v_{xxx}\right)\\
&\quad + \left(\frac{(\gamma-1)}{-4\beta\lambda+\gamma^2-\gamma} - \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_{xx} - tu_{xy}\right)\left(\beta v u_x^2 - v u_y + v\delta - \alpha v_{xx}\right)\\
&\quad - \left(\frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_{xxx} + tu_{xxy}\right)\alpha v_x + \left(\frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_{xxxx} + tu_{xxxy}\right)\alpha v,\\
C^3 &= -tv\left(u_{xt} - \alpha u_{xxxx} + \beta u_x^2 u_{xx} - u_{xx}u_y + \gamma u_x u_{xy} + \lambda u_{yy} + \delta u_{xx}\right)\\
&\quad + \left(\frac{(\gamma-1)x}{-4\beta\lambda+\gamma^2-\gamma} - \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_x - tu_y\right)\left(-v u_{xx} - \gamma v_x u_x - \gamma v u_{xx} - \lambda v_y\right)\\
&\quad + \left(\frac{(\gamma-1)}{-4\beta\lambda+\gamma^2-\gamma} - \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_{xx} - tu_{xy}\right)\gamma v u_x - \left(\frac{2\beta}{-4\beta\lambda+\gamma^2-\gamma}u_x + \frac{2\beta y}{-4\beta\lambda+\gamma^2-\gamma}u_{xy} + tu_{yy}\right)\lambda v.
\end{aligned} \tag{74}$$

This vector involves an arbitrary solution v of the adjoint equation Eq.(66) and provides an infinite number of the conservation laws.

Remark We have checked all the exact solutions satisfy the original Eq.(1) , and the above solutions of the extended (2+1)-dimensional Jaulent-Miodek equation have not been found in other references. To our knowledge, the symmetry reductions obtained in this paper are completely new and has not been studied yet in previous papers.

5. conclusion

In this paper, we investigated the extended (2+1)-dimensional Jaulent-Miodek equation via Lie symmetries. By applying the direct symmetry method ,we obtain the classical Lie point symmetry of the equation. Based on given Lie symmetry , we reduce the extended (2+1)-dimensional Jaulent-Miodek equation to the (1+1)- dimensional PDE and ODE. We also have derived some exact solutions of Eq.(1) by using the relationship between the new solutions and the old ones. In order to reflect the characteristics and properties of this solutions, we give figures of some solutions. At last, we give the conservation laws of Eq.(1). These conclusions may be useful for the explanation of some practical physical problems.

The symmetry study plays a very important role in almost all scientific fields, especially in the soliton theory because of the existence of infinitely many symmetries for integrable system. Furthermore, the symmetry analysis based on the Lie group method is a very powerful method and is worthy of being studyied further.

6. Acknowledgments

This work is supported by National Natural Science Foundation of China under Grant(Nos.11505090), Research Award Foundation for Outstanding Young Scientists of Shandong Province(No.BS2015SF009) .

References

References

[1] G.W. Bluman and S. Kumei, Symmetries and Differential Equations, Springer-Verlag, New York, 1989.

[2] S. Lie, Über die Integration durch bestimmte Integrale von einer Klasse linearer partieller Differential gleichungen, Arch. Math. 328 (1881).

[3] H.Z. Liu, J.B. Li, Symmetry reductions, dynamical behavior and exact explicit solutions to the Gordon types of equations, Appl. Math. Comput. 257 (2014) 144-156.

[4] J.H. He, X.H. Wu, Exp-function method for nonlinear wave equation, Chaos, Solitions and Fractals. 3(2006) 700-708.

[5] E.J. Parkes, B.R. Duffy, An automated tanh-function method for finding solitary wave solutions to non-linear evolution equations, Comput. Phys. Commun. 98(1996) 288-300.

[6] A.M. Wazwaz, The extended tanh method for the ZakharovCKuznetsov (ZK) equation, the modified ZK equation, and its generalized forms, Commun. Nonlinear. Sci. Numer. Simul. 13(2008) 1039-1047.

[7] X.Q. Zhao, H.Y. Zhi, H.Q. Zhang, Improved Jacobi-function method with symbolic computation to construct new double-periodic solutions for the generalized Ito system, Chaos, Solitions and Fractals. 28(2006) 112-126.

[8] H.Q. Zhang, Extended Jacobi elliptic function expansion method and its applications, Commun. Nonlinear. Sci. Numer. Simul. 12(2007) 627-635.

[9] J.F. Zhang, C.Q. Dai, Q.Yang, J.M. Zhu, Variable-coefficient F-expansion method and its application to nonlinear Schrödinger equation, Opt. Commun. 252(2005) 408-421.

[10] R. Hirota, J. Satsuma, Soliton solutions of a coupled KortewegCde Vries equation, Phys. Lett. A. 85(1981) 407-408.

[11] R. Conte, M. Musette, Painleve analysis and Backlund transformation in the KuramotoCSivashinsky equation, J. Phys. A: Math. Gen. 22(1989) 169.

[12] Y. Matsuno, Reduction of dispersionless coupled KdV equations to the Euler-Darboux eqution, J. Math. Phys. 42(2001) 1744-1760.

[13] H.Y. Ruan, S.Y. Lou, New symmetries of the JaulentłMiodek hierarchy, J. Phys. Soc. Jpn. 62(1993) 1917-1921.

[14] Y.B. Zeng, Separability and dynamical r-matrix for the constrained flows of the Jaulent-Miodek hiodek hierarchy, Phys. Lett. A. 216(1996) 26-32.

[15] E.G. Fan, Uniformly constructing a series of explicit exact solutions to nonlinear equations mathematical physics, Chaos, Solitions and Fractals. 16(2003) 819-839.

[16] D.H. Feng, J.B. Li, Bifurcations of travelling wave solutions for Jaulent-Miodek equations, Appl. Math. Mech. -Engl. Ed. 28(2007) 999-1005.

[17] A.M. Wazwaz, The tanh-coth and the sech methods for exact solutions of the Jaulent-Miodek equation, Phys. Lett. A. 366(2007) 85-90.

[18] J.H. He, L.N. Zhang, Generalized solitary solution and compacton-like solution of the Jaulent-Miodek equations using the Exp-function method, Phys. Lett. A. 372(2008) 1044-1047.

[19] X.H. Du, Using Trial Equation Method To Solve New Exact Traveling Wave Solutions to JaulentłMiodek Equation, Math. Practice. Theory. 6(2010) 204-208.

[20] Kavitha L, Sathishkumar P, Nathiyaa T, et al., Cusp-like singular soliton solutions of Jaulent-Miodek equation using symbolic computation, Physica Scripta. 79(2009) 035403.

[21] A.M. Wazwaz, Multiple kink solutions and multiple singular kink solutions for (2 + 1)-dimensional nonlinear models generated by the Jaulent-Miodek hierarchy, Phys. Lett. A. 373 (2009) 1844-1846.

[22] Y.Y. Zhang, X.Q. Liu, Symmetry reductions and exact solutions of the (2 + 1)-dimensional JaulentCMiodek equation, Appl. Math. Comput. 219(2012) 911-916.

[23] X.P. Xin, X.Q. Liu, L.L. Zhang, Explicit solutions of the Bogoyavlensky-Konoplechenko equation, Appl. Math. Comput. 215(2010) 3669-3673.

[24] Y.Y. Zhang, G.W. Wang, X.Q. Liu, Symmetry reductions and explicit solution of the (2+1)-dimensional nonlinear evolution, Chin. J. Quantum. Elect. 29(2012) 411-416.